

A-Level Edexcel

# PHYSICS

UNIT 5(IAL)

2020 — 2025

Chapter 1	<b>Mechanics</b>	Page 1
Chapter 2	<b>Materials</b>	Page 3
Chapter 3	<b>Waves and Particle Nature of Light</b>	Page 4
Chapter 4	<b>Electric Circuits</b>	-----
Chapter 5	<b>Further Mechanics</b>	Page 6
Chapter 6	<b>Electric and Magnetic Fields</b>	Page 9
Chapter 7	<b>Nuclear and Particle Physics</b>	Page 10
Chapter 8	<b>Thermodynamics</b>	Page 18
Chapter 9	<b>Nuclear Decay</b>	Page 59
Chapter 10	<b>Oscillations</b>	Page 104
Chapter 11	<b>Astrophysics and Cosmology</b>	Page 151
	<b>Answers</b>	Page 236

1 - (WPH11/5(IAL)\_Summer\_2024\_Q12) - *Mechanics, Astrophysics And Cosmology*

The planet Uranus has nearly thirty moons. Titania is the largest of these moons. The radius of Titania is  $7.88 \times 10^5$  m. The mass of Titania is  $3.40 \times 10^{21}$  kg.

Scientists plan to use a vehicle with a mass of 210 kg to explore Titania.

Calculate the weight of this vehicle on the surface of Titania.

Weight = .....

2 - (WPH11/5(IAL)\_Summer\_2024\_Q13) - *Thermodynamics, Mechanics*

The photograph shows an airship. This consists of a large balloon filled with helium gas with a basket suspended beneath.



(Source: © Tom Bushey/Alamy Stock Photo)

(a) Explain how the airship is able to float in the air.

(2)

(b) The pressure of helium inside the balloon is  $1.08 \times 10^5$  Pa at a temperature of  $25^\circ\text{C}$ .

Calculate the mass of helium gas in the balloon.

volume of balloon =  $7020 \text{ m}^3$

mass of a helium atom =  $6.64 \times 10^{-27} \text{ kg}$

(4)

Mass of helium gas = .....

1 - (WPH11/5(IAL)\_Winter\_2025\_Q2) - *Materials, Oscillations*

Ductile materials can be used in buildings to reduce the amplitude of forced oscillations during an earthquake.

Which of the following is the reason that ductile materials are used?

- A These materials are stiff.
- B These materials are strong.
- C These materials deform elastically.
- D These materials deform plastically.

www.exam-mate.com

**1** - (WPH11/5(IAL)\_Summer\_2024\_Q16) - *Astrophysics And Cosmology, Waves And Particle Nature Of Light*

The star  $\epsilon$ -Eridani is one of the closest stars to the Sun. The intensity of  $\epsilon$ -Eridani, as measured on the Earth, is  $1.05 \times 10^{-9} \text{ W m}^{-2}$ .

- (a) Calculate the wavelength,  $\lambda_{\text{max}}$ , corresponding to the peak intensity of emission of radiation from  $\epsilon$ -Eridani.

distance of  $\epsilon$ -Eridani from the Earth =  $9.94 \times 10^{16} \text{ m}$   
radius of  $\epsilon$ -Eridani =  $5.12 \times 10^8 \text{ m}$

(5)

$$\lambda_{\text{max}} = \dots\dots\dots$$

- (b)  $\epsilon$ -Eridani is moving away from the Earth with a velocity of  $1.55 \times 10^4 \text{ m s}^{-1}$ .

A scientist observed the light received from  $\epsilon$ -Eridani using a spectrometer.


Deduce whether the scientist was able to detect a change in the wavelength of light, of wavelength 480 nm, emitted by the star.

resolution of spectrometer = 0.05 nm

(3)

2 - (WPH11/5(IAL)\_Winter\_2025\_Q18) - Astrophysics And Cosmology, Waves And Particle Nature Of Light

The diagram shows dark lines in the continuous spectrum of electromagnetic radiation from the Sun.




- (a) Atoms in the outer regions of the Sun absorb electromagnetic radiation. This produces the dark lines in the spectrum.

Explain why electromagnetic radiation is only absorbed at some wavelengths.

(3)

- (b) The Sun rotates on its axis. The diagram shows the Sun rotating at its 'equator'.



(Source: © NASA / SDO / SCIENCE PHOTO LIBRARY)

One of the dark lines in the spectrum occurs at a wavelength of 589 nm.

Radiation is received on Earth from opposite ends of the Sun's diameter. There is a difference between the wavelengths of radiation received from opposite ends due to the Sun rotating about its axis.

The difference in wavelength is  $7.94 \times 10^{-3}$  nm.

A website states that at its equator, the Sun rotates once every 27 days about its axis.

Assess the accuracy of this statement.

radius of Sun =  $6.96 \times 10^8$  m

1 day =  $8.64 \times 10^4$  s

(5)

# ANSWERS

[www.exam-mate.com](http://www.exam-mate.com)

## 1 - (WPH11/5(IAL)\_Summer\_2024\_Q12) - Mechanics, Astrophysics And Cosmology

	Use of $g = \frac{GM}{r^2}$	(1)	3
	Use of $W = mg$	(1)	
	$W = 77 \text{ N}$	(1)	
	<u>Example of calculation</u> $g = \frac{6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2} \times 3.40 \times 10^{21} \text{ kg}}{(7.88 \times 10^5 \text{ m})^2} = 0.365 \text{ N kg}^{-1}$ $F = 210 \text{ kg} \times 0.365 \text{ N kg}^{-1} = 76.7 \text{ N}$		

## 2 - (WPH11/5(IAL)\_Summer\_2024\_Q13) - Thermodynamics, Mechanics

(a)	<u>Upthrust</u> on airship is equal to the <u>weight</u> of the airship	(1)	2
	(so) resultant force on the airship is zero (so airship floats in the air) <b>Or</b> (vertically) balanced forces act on the airship	(1)	
	If neither MP seen, max 1 mark for statement indicating that upwards force is equal to downwards force.		
(b)	Use of $pV = NkT$	(1)	4
	Conversion of temperature to K	(1)	
	Use of $M = Nm$	(1)	
	$M = 1200 \text{ kg}$	(1)	
	<u>Example of calculation</u> $N = \frac{pV}{kT} = \frac{1.08 \times 10^5 \text{ Pa} \times 7020 \text{ m}^3}{1.38 \times 10^{-23} \text{ m}^2 \text{ kg s}^{-2} \text{ K}^{-1} \times (273 + 25) \text{ K}} = 1.84 \times 10^{29}$ $M = 1.84 \times 10^{29} \times 6.64 \times 10^{-27} \text{ kg} = 1220 \text{ kg}$		

1 - (WPH11/5(IAL)\_Winter\_2025\_Q2) - *Materials, Oscillations*

	<p><b>D is the correct answer (these materials deform plastically)</b></p> <p>A is not correct because this will not absorb energy from the oscillation B is not correct because this will not absorb energy from the oscillation C is not correct because this will absorb and then return energy to the oscillation</p>	<p><b>1</b></p>
--	---	-----------------

1 - (WPH11/5(IAL)\_Summer\_2024\_Q16) - Astrophysics And Cosmology, Waves And Particle Nature Of Light

<b>(a)</b>	Use of $I = \frac{L}{4\pi d^2}$ (1)	5
	Use of $L = \sigma AT^4$ (1)	
	Use of $A = 4\pi r^2$ (1)	
	Use of $\lambda_{\max} T = 2.898 \times 10^{-3} \text{ m K}$ (1)	
	$\lambda_{\max} = 5.64 \times 10^{-7} \text{ m}$ (1)	
	<p><u>Example of calculation</u></p> $1.05 \times 10^{-9} \text{ W m}^{-2} = \frac{L}{4\pi \times (9.94 \times 10^{16} \text{ m})^2}$ $L = 4\pi \times (9.94 \times 10^{16} \text{ m})^2 \times 1.05 \times 10^{-9} \text{ W m}^{-2} = 1.30 \times 10^{26} \text{ W}$ $A = 4\pi \times (5.12 \times 10^8 \text{ m})^2 = 3.29 \times 10^{18} \text{ m}^2$ $1.30 \times 10^{26} \text{ W} = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{K}^{-4} \times 3.29 \times 10^{18} \text{ m}^2 \times T^4$ $T = \sqrt[4]{\frac{1.30 \times 10^{26} \text{ W}}{5.67 \times 10^{-8} \text{ W m}^{-2} \text{K}^{-4} \times 3.29 \times 10^{18} \text{ m}^2}} = 5140 \text{ K}$ $\lambda_{\max} = \frac{2.898 \times 10^{-3} \text{ m K}}{5140 \text{ K}} = 5.64 \times 10^{-7} \text{ m}$	
<b>(b)</b>	Use of $\frac{\Delta\lambda}{\lambda} = \frac{v}{c}$ (1) [Allow a 2 step approach using a calculation of the red shift, z ]	3
$\Delta\lambda = 0.025 \text{ nm}$ (1)		
0.025 (nm) < 0.05 (nm), so no wavelength change could be detected (1)		
<b>Or</b> comparison of calculated value with 0.05 (nm) and consistent conclusion made. (1)		
<p><u>Example of calculation</u></p> $\frac{\Delta\lambda}{480 \text{ nm}} = \frac{1.55 \times 10^4 \text{ m s}^{-1}}{3.00 \times 10^8 \text{ m s}^{-1}}$ $\Delta\lambda = 480 \text{ nm} \times 5.17 \times 10^{-5} = 0.0248 \text{ nm}$		

2 - (WPH11/5(IAL)\_Winter\_2025\_Q18) - Astrophysics And Cosmology, Waves And Particle Nature Of Light

<b>(a)</b>	<p>Electron/atomic energy levels are discrete  <b>Or</b> Electrons can only have certain energy states (1)</p> <p>A <u>photon</u> is absorbed with an energy equal to the difference in the energy between two energy levels (1)</p> <p>Only certain (energy level) transitions are possible (so only some wavelengths/frequencies are absorbed) (1)</p>	<b>3</b>
<b>(b)</b>	<p>Use of <math>\frac{\Delta\lambda}{\lambda} \approx \frac{v}{c}</math> (1)</p> <p>Use of <math>v = \omega r</math> (1)</p> <p>Use of <math>\omega = \frac{2\pi}{T}</math> (1)</p> <p>Conversion between seconds and days (1)</p> <p><math>T = 25 \text{ days} &lt; 27 \text{ days}</math>, so statement not accurate (1)</p> <p><b>Or</b> <math>T = 25 \text{ days} \approx 27 \text{ days}</math>, so statement accurate</p> <p><b>ALTERNATIVE SOLUTION:</b></p> <p>Use of <math>\omega = \frac{2\pi}{T}</math> to calculate angular velocity of Sun (1)</p> <p>Use of <math>v = \omega r</math> to calculate rotational velocity of (edge of) Sun (1)</p> <p>Use of <math>\frac{\Delta\lambda}{\lambda} \approx \frac{v}{c}</math> to calculate wavelength for approaching and receding side (1)</p> <p>Calculate difference in <math>\lambda</math> for each side of Sun (1)</p> <p><math>\Delta\lambda = 7.36 \times 10^{-3} \text{ nm} &lt; 7.94 \times 10^{-3} \text{ nm}</math>, so statement is inaccurate (1)</p> <p><u>Example of calculation</u></p> $\Delta\lambda = \frac{7.94 \times 10^{-3} \text{ nm}}{2} = 3.97 \times 10^{-3} \text{ nm}$ $v = \left( \frac{3.97 \times 10^{-3} \text{ nm}}{589 \text{ nm}} \right) \times 3.00 \times 10^8 \text{ m s}^{-1} = 2.02 \times 10^3 \text{ m s}^{-1}$ $\omega = \frac{2.02 \times 10^3 \text{ m s}^{-1}}{6.96 \times 10^8 \text{ m}} = 2.91 \times 10^{-6} \text{ rad s}^{-1}$ $T = \frac{2\pi \text{ rad}}{2.91 \times 10^{-6} \text{ rad s}^{-1}} = 2.16 \times 10^6 \text{ s}$ $T = \frac{2.16 \times 10^6 \text{ s}}{8.64 \times 10^4 \text{ s day}^{-1}} = 25 \text{ days}$	<b>5</b>

1 - (WPH11/5(IAL)\_Summer\_2024\_Q18) - Astrophysics And Cosmology, Further Mechanics

<b>(a)</b>	Gravitational force equated to centripetal force (1)  Substitution of $\omega = \frac{2\pi}{T}$ (1) <b>Or</b> Substitution of $v = \frac{2\pi r}{T}$ (1)  Algebra to obtain required expression (1)  <u>Example of derivation</u> $m\omega^2 r = \frac{GMm}{r^2}$ $m \left( \frac{2\pi}{T} \right)^2 r = \frac{GMm}{r^2}$ $T^2 = \frac{4\pi^2}{GM} r^3$	(1)  (1)  (1)  <b>3</b>
<b>(b)(i)</b>	Use of $T^2 = \frac{4\pi^2 r^3}{GM}$ (1)  $T = 8.3 \times 10^{15}$ (s) (1)  <u>Example of calculation</u>  $T = \sqrt{\frac{4\pi^2 \times (5.7 \times 10^{20} \text{ m})^3}{6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2} \times 8.0 \times 10^{11} \times 1.99 \times 10^{30} \text{ kg}}}$ $\therefore T = 8.30 \times 10^{15} \text{ s}$	(1)  (1)  <b>2</b>
<b>(b)(ii)</b>	The actual period is (much) smaller than the calculated value (1)  So the mass of the galaxy must be greater than $8.0 \times 10^{11}$ solar masses [accept the given mass for the numerical value] <b>Or</b> the gravitational force on the star must be bigger (than assumed) (1)  There must be matter that does not emit em-radiation <b>Or</b> There must be matter that we cannot detect via em-radiation [Accept "there must be matter that we cannot see"] (1)  (This suggests that) there is dark matter (1)	(1)  (1)  (1)  (1)  <b>4</b>

## 2 - (WPH11/5(IAL)\_Summer\_2025\_Q17) - Thermodynamics, Further Mechanics, Astrophysics And Cosmology

(a)	Substitute into $V = \frac{4}{3}\pi r^3$ Use of $pV = NkT$ Conversion of temperature to kelvin Use of number of molecules to calculate mass of gas $m = 230 \text{ kg}$	(1) (1) (1) (1) (1)	Do not credit $pV = nRT$ unless final answer is correct If answer incorrect, <i>could</i> award marks for MP1, MP3, MP4  <u>Example of calculation</u> $V = \frac{4}{3}\pi \times (3.5 \text{ m})^3 = 179.6 \text{ m}^3$ $N = \frac{1.12 \times 10^5 \text{ Pa} \times 179.6 \text{ m}^3}{1.38 \times 10^{-23} \text{ J K}^{-1} \times (273 + 22) \text{ K}} = 4.94 \times 10^{27}$ $m = 4.94 \times 10^{27} \times 4.67 \times 10^{-26} \text{ kg} = 231 \text{ kg}$	5
(b)	Use of $\omega = \frac{2\pi}{T}$ $T = 228 \text{ s}$	(1) (1)	<u>Example of calculation</u> $T = \frac{2\pi \text{ rad}}{2.76 \times 10^{-2} \text{ rad s}^{-1}} = 227.6 \text{ s}$	2
(c)	Use of $g = \frac{GM}{r^2}$ $M = 1.8 \times 10^{12} \text{ kg}$	(1) (1)	<u>Example of calculation</u> $M = \frac{gr^2}{G} = \frac{9.8 \text{ N kg}^{-1} \times (3.5 \text{ m})^2}{6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}} = 1.80 \times 10^{12} \text{ kg}$	2

## 3 - (WPH11/5(IAL)\_Winter\_2025\_Q3) - Further Mechanics, Electric And Magnetic Fields

	<b>C is the correct answer (both fields can produce repulsive forces)</b>  A is not correct because both fields are radial B is not correct because both fields can exert attractive forces D is not correct because both fields obey an inverse square law for force	1
--	---	---